

2

AD-A201 242

DTIC FILE COPY

OFFICE OF NAVAL RESEARCH
PROGRAM ELEMENT 61153N
TECHNICAL REPORT NO. 6

NRL BORON PROPELLANTS COMBUSTION PROGRAM

BY

H.H. NELSON

Published

in the

Proceedings of the 1988 AFOSR/ONR
Contractor's Meeting on Combustion
15-19 June 1988, Los Angeles, CA

DTIC
ELECTE
OCT 25 1988
S D

CHEMISTRY DIVISION
NAVAL RESEARCH LABORATORY
WASHINGTON, DC 20375-5000

JUNE 1988

Reproduction in whole or in part is permitted for
any purpose of the United States Government

This document has been approved for public release
and sale, its distribution is unlimited

88 10 24 019

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b. RESTRICTIVE MARKINGS NONE		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION / AVAILABILITY OF REPORT UNLIMITED		
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S)			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Naval Research Laboratory		6b. OFFICE SYMBOL (if applicable)	7a. NAME OF MONITORING ORGANIZATION Chemistry Division Office of the Chief of Naval Research		
6c. ADDRESS (City, State, and ZIP Code) Washington, DC 20375-5000			7b. ADDRESS (City, State, and ZIP Code)		
8a. NAME OF FUNDING / SPONSORING ORGANIZATION ONR		8b. OFFICE SYMBOL (if applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER		
8c. ADDRESS (City, State, and ZIP Code)			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO. 61153N	PROJECT NO. RR013-01-4C	TASK NO.
11. TITLE (Include Security Classification) NRL Boron Propellants Combustion Program(U)					
12. PERSONAL AUTHOR(S) H.H. Nelson					
13a. TYPE OF REPORT		13b. TIME COVERED FROM 9/87 TO 9/88		14. DATE OF REPORT (Year, Month, Day) JUNE 1988	
15. PAGE COUNT					
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP			
19. ABSTRACT (Continue on reverse if necessary and identify by block number) The NRL program on High-Energy Propellant Chemistry includes work in three project areas. The first of these is a mass spectrometry based study of the oxide layer formed at the surface of elemental boron. The second involves the study of chemical mechanisms of homogeneous boron combustion. This work has focussed on boron hydrides in the past year but is expanding to cover boron oxyhydrides. The third area is a theoretical investigation of important intermediates in boron combustion. Structures, thermodynamics and reaction pathways are being studied. All three projects are oriented to assist in the understanding and control of heterogeneous and homogeneous combustion of boron and boron-doped fuels.					
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED		
22a. NAME OF RESPONSIBLE INDIVIDUAL J.R. McDonald			22b. TELEPHONE (Include Area Code) (202) 767-3340		22c. OFFICE SYMBOL

NRL BORON PROPELLANTS COMBUSTION PROGRAM

H.H. Nelson
Chemistry Division/Code 6110
Naval Research Laboratory
Washington, DC 20375-5000

SUMMARY/OVERVIEW:

The NRL program on High-Energy Propellant Chemistry includes work in three project areas. The first of these is a mass spectrometry based study of the oxide layer formed at the surface of elemental boron. The second involves the study of chemical mechanisms of homogeneous boron combustion. This work has focussed on boron hydrides in the past year but is expanding to cover boron oxyhydrides. The third area is a theoretical investigation of important intermediates in boron combustion. Structures, thermodynamics and reaction pathways are being studied. All three projects are oriented to assist in the understanding and control of heterogeneous and homogeneous combustion of boron and boron-doped fuels.

TECHNICAL DISCUSSION:

The oxide coating that forms rapidly on boron (B_2O_3)_n has been investigated by fast-atom bombardment mass spectrometry. Bombardment of the glassy surface of oxidized boron by high energy xenon atoms yielded twenty-two gas-phase boron oxide ions that were heretofore unknown.[1,2] Collisional activation of these ions has revealed structural elements that are common to the most stable boron oxide ions. These ions exist with boron limited to two- or three-coordination with oxygen and are thus built upon integral BO_2 triangles and terminal $-B=O$ units.

We have studied the reactions of BH_3 [3] and BH with various oxidants and hydrocarbons. The BH_3 radical is produced by 193nm photolysis of B_2H_6 and monitored by infrared diode laser absorption near 1140 cm^{-1} . The BH is produced by BH_3CO photolysis and probed by UV lif near 430 nm. We have measured rate constants for the reactions of BH_3 with CO , NO and C_2H_4 and upper limits to the rate constants for reaction with O_2 and H_2O . Rate constants for reaction of BH with NO , H_2O , O_2 , CO_2 , C_2H_4 , TME , CO and H_2 have also been measured and will be discussed in the presentation.

Finally, we have been performing ab-initio quantum chemical calculations of potential energy surfaces for species important to boron combustion. Efforts have focussed on the theoretical prediction of heats of formation for small boron hydrides and a few oxygen containing species and on kinetic studies involving critical species.

We have recently completed a theoretical study of the reaction $H_2 + BO \rightarrow H + HBO$. Our calculations predict this reaction to have a zero point corrected barrier of 9.5 kcal/mole and an exoergicity of 6.4 kcal/mole. The calculated rate constant is well represented over a wide temperature range by the three parameter expression:

$$k(T) = 2.96 \times 10^{-22} T^{3.29} e^{-4.40/RT}$$

We are currently performing a study of the reaction between BO and water. The potential energy

surface calculations are done using multiconfiguration SCF and multireference CI techniques. The rates are computed using conventional or variational transition state theory with quantum corrections along the reaction coordinate. Also underway is the calculation of the regions of the BH_3 potential energy surface important to the unimolecular dissociation to BH and molecular hydrogen.

REFERENCES

1. Doyle, R.J., Jr., Anal. Chem. 1987, 59, 537-539.
2. Doyle, R.J., Jr., J. Am. Chem. Soc. In Press, June 1988 publication.
3. L. Pasternack, R.J. Balla and H.H. Nelson, J. Phys. Chem. 92, 1200 (1988).

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution	
Availability Codes	
Dist	Availability Codes
A-1	

